

Cardiac Functions & Lipid Profile in Obese Children & Adolescents

Dr .T. Ajith Kumar¹, Dr. Vineeta Pande², Dr. Sharad Agarkhedkar³, Dr. Mayank Surana¹

¹Resident, ²Professor, ³Professor & Head, Department of Paediatrics, Dr. D. Y. Patil Vidyapeeth's (DPU) Dr. D. Y. Patil Medical College & Hospital, Pune- 411018

Corresponding Author : Dr. Vineeta Pande

E-mail : ajithkumar127@gmail.com

Address : Department of Paediatrics, Dr. D. Y. Patil Vidyapeeth's (DPU) Dr. D. Y. Patil Medical College & Hospital, Pune- 411018

Abstract :

Background: Obesity is a disease which results from genetic or lifestyle factors. Such diseases are called Non communicable diseases. The epidemic of obesity among youth is spreading at an alarming rate due to lack of physical activity, dietary habits. The percentage of youths who are at risk of becoming overweight continues to increase. National data indicates that 16% of children aged six to nineteen years are overweight.⁽¹⁾ Considering the high prevalence of obesity among children and adolescents and its associated cardiovascular complications like hypertension, left ventricular hypertrophy, increased left ventricular mass, decreased left ventricular ejection fraction, the present study was planned to assess cardiac functions and lipid profile in obese children and adolescents. Early detection and intervention can help in reducing the complications associated with obesity. **Objective:** Primary objective is to assess cardiac functions and lipid profile in obese children and adolescents. **Methodology:** 100 obese children and adolescents in the age group 6-18 years were taken after excluding obese children with evidence of endocrine disease, malformation syndromes and iatrogenic obesity (drug treatments). **Results:** In the present study 7% subjects were hypertensive while 8% were prehypertensive. 71.43% children with hypertension were performing physical activity for < 30 min in 1-3 days/week while 49.11% normotensive children were performing physical activity for > 30 min per day. 3% obese children were having abnormal LVEF on 2DECHO .All the 3 obese children with abnormal LVEF were having physical activity less than 30 min in 1-3 days /week and the difference was statistically significant. All the three

children with abnormal LVEF were hypertensive and the difference was statistically significant. 6% children were having raised serum cholesterol >190mg/dl. Serum triglyceride >150mg/dl was observed in 17% obese children. HDL level <20mg/dl was observed in 7% children while LDL above 130mg/dl was observed in 6% children. The proportion among male and female children was comparable. **Conclusion:** Cardiovascular complications of adulthood in obese children and adolescents may be prevented by early identification and intervention in the form of healthy life style, dietary intake and physical activity.

Keywords: Obesity, Cardiac functions, Lipid profile.

Introduction:

The problem of overweight and obesity can be seen among children and adolescents of developed as well as developing countries. Previously obesity was exclusively seen in adults, becoming more prevalent with advancing age, however with children's sedentary lifestyles due to a number of factors, over consumption of high-caloric foods, excessive TV viewing during meals, obesity in children is also on the rise. Children with high cholesterol level were found to suffer from increased cholesterol and cardiovascular complications in adulthood.⁽²⁾ The cardiovascular complications associated with obesity include hypertension, left ventricular hypertrophy, increased left ventricular mass, decreased left ventricular ejection fraction. These children are at high risk for psychological and social adjustment problem, low self-esteem, symptoms of depression and increased suicidal thoughts and attempts. Musculoskeletal symptoms include impaired mobility and fractures. Life style modifications and adequate physical activity play a major role in preventing, controlling obesity and its associated complications.

Methodology:

Cross sectional observational study conducted in 100 obese children and adolescents in the age group of 6-18 years attending paediatric OPD of tertiary care hospital from August 2017-August 2019. Obese children with evidence of endocrine disease, malformation syndromes and iatrogenic obesity (drug treatments) were excluded. After obtaining the consent, history was taken, required physical examination and necessary investigations were done as per proforma. Blood pressure was recorded in supine position on three different occasions and compared with IAP BP centile charts.

Anthropometric measurements including weight, height, waist circumference, BMI was done. Lipid profile was assessed with early morning blood samples taken after a minimum of 12 hour fasting. Cardiac functions were assessed by 2D ECHO. All the data collected was entered in MS EXCEL and analysed by using SPSS. Chi square test and ANOVA test was applied to check the statistical association.

Results:

Table 1: Distribution of study population according to Age groups, Gender & Socioeconomic status

Character	Age Groups (Years)			Gender		Socioeconomic status		
	6-8	9-11	12-18	Boys	Girls	Upper middle class	Upper lower class	Lower middle class
Percentage	29	44	27	57	43	51	31	18

Table 2: Distribution of cases according to physical activity in study group

Physical activity		No of cases	Percentage (%)
Daily (n=61)	>30 min	43	43
	<30 min	18	18
1 – 3 day/ week (n=39)	>30 min	26	26
	<30 min	13	13
Total		100	100

Table 3: Association between physical activity and blood pressure in study group

Physical activity		Blood pressure			Total
		Hyper tension	Prehyper tension	Normo tension	
Daily (n=61)	>30 min	0	1	42	43
	<30 min	2	4	12	18
1 – 3 day /week (n=39)	>30 min	0	0	26	26
	<30 min	5	3	5	13
Total		7	8	85	100

Chi-square = 41.23, P<0.0001

Table 4: Lipid profile distribution of cases according to gender in study group

Lipid profile		Male (n=57)	Female (n=43)	Total (n=100)
Sr. Cholesterol (mg/dl)	>190	4	2	6
	<190	53	41	94
Sr. Triglyceride (mg/dl)	>150	12	5	17
	<150	45	38	83
HDL (mg/dl)	<20	2	5	7
	>20	55	38	93
LDL (mg/dl)	>130	5	1	6
	<130	52	42	94

Table 5: 2D ECHO finding distribution of cases in study group

2D ECHO finding	No of cases	Percentage
Abnormal LVEF	3	3
Normal LVEF	97	97
Total	100	100.0

Table 6: Association between blood pressure and 2D ECHO finding in study group

Blood pressure	2D ECHO finding		Total
	Abnormal	Normal	
Hypertension	3	4	7
Prehypertension	0	8	8
Normotension	0	85	85
Total	3	97	100

Chi-square = 41.09, P<0.0001

Table 7: Association between physical activity and 2D ECHO finding in study group

Physical activity		2D ECHO finding		Total
		Abnormal	Normal	
Daily (n=61)	>30 min	0	43	43
	<30 min	0	18	18
1 – 3 day/ week (n=39)	>30 min	0	26	26
	<30 min	3	10	13
Total		3	97	100

Chi-square = 20.7, P<0.0001

Discussion:

In the present study it was seen that majority of the children were in the age group of 9-11 years (44%) followed by 6-8 years (29%), 12-14 years (25%) and 15 – 18 years (2%). In Nascimento H⁽³⁾ study mean age of subjects was 10.8 ± 3.0 yrs while in A.M. Zoair *et al*⁽⁴⁾ mean age of obese group was 10.56 ± 2.82 yrs.

The present study observed that majority of the obese children were male (57%) followed by female were observed (43%). In the Minakshi B⁽⁵⁾ study males were 52% cases and females were 48% cases. LaFrance *et al*⁽⁶⁾ stated that boys' obesity was higher than girls' obesity. In the present study it was observed that 51% children in the study were belonging to upper middle class while 31% were from upper lower class and 18% were from lower middle class. Shabana *et al*⁽⁷⁾ also observed higher prevalence of obesity among Upper Socio Economic class as compared to those among children of Lower Socio Economic Class.

In the present study it was noticed that 43% children were involved in daily physical activity for more than 30min while 26% were doing physical activity 1-3days/week for more than 30min (**discussed in table 2**). Similar results are shown in the study done by S. Kumar *et al*⁽⁸⁾ in Davengere which showed that those with absent physical activity had 2 times more risk of falling in obese group as compared to those who had more physical activity ($p < 0.001$). The present study observed that out of total 100 obese children 7% were hypertensive while 8% were pre hypertensive and remaining 85% were normotensive. In the present study it was seen that 71.43% children with hypertension were performing physical activity for < 30 min in 1-3 days/week while 49.11% normotensive children were performing physical activity for > 30 min per day (**discussed in table 3**). In this study we observed that 6% children were having raised serum cholesterol > 190 mg/dl. Serum triglyceride > 150 mg/dl was

observed in 17% obese children. HDL level < 20 mg/dl was observed in 7% children while LDL above 130mg/dl was observed in 6% children. The proportion among male and female children was comparable. Higher serum cholesterol, serum triglyceride, serum LDL levels are seen in boys while lower serum HDL levels are seen in girls (**discussed in table 4**). Similarly in Minakshi B⁽⁵⁾ study 85% cases had TGL levels less than 150 mg/dl and 15% had high or equal to 150mg/dl while 69% of the cases had HDL-C less than 40mg/dl and 31% of the cases had HDL-C more than or equal to 40mg/dl. This was in accordance with Freedman *et al*⁽⁹⁾ and Reinhret *et al*⁽¹⁰⁾ who found that overweight is strongly related to concentrations of HDL-cholesterol and TG and weakly related to concentrations of LDL-cholesterol; while Kim *et al*⁽¹¹⁾ found that all types of lipids were significantly associated with overweight and obesity. Also, Chinali *et al*⁽¹²⁾ stated that obese adolescents had significantly higher total TG and lower high density lipoprotein.

In our study 3% obese children were having abnormal LVEF on 2D ECHO findings while remaining 97% were having normal LVEF (discussed in table 5). Obese adolescents exhibit changes in the left ventricular (LV) mass related to increase in cardiac workload. All the three children with abnormal LVEF on 2D ECHO were hypertensive and the difference was statistically significant (**discussed in table 6**). In the present study it was seen that all the 3 obese children with abnormal LVEF on 2D ECHO were having physical activity less than 30 min in 1-3 days /week and the difference was statistically significant (**discussed in table 7**). Antonio Vitarelli *et al*⁽¹³⁾ assessed the presence of early myocardial deformation abnormalities in dyslipidemic children free from other cardiovascular risk factors, using 2-dimensional speckle tracking echocardiography (2DSTE) and 3-dimensional speckle tracking echocardiography (3DSTE).

Conclusion:

Cardiac functions were abnormal in 3% of obese children and adolescents. Abnormal LVEF on 2D ECHO finding were associated with less physical activity and Hypertension with statistically significant difference, while no significant association was observed between lipid profile and echo findings among obese children and adolescents. Cardiovascular complications of adulthood in obese children and adolescents may be prevented by early identification and intervention in the form of healthy life style, dietary intake and physical activity.

References:

1. Hedly, (2004). Nutrition for the preschool child. In pediatric manual of Clinical dietetics, ed 2. Chicago, IL: American dietetic association, 77 - 82.
2. Shen GX. Oxidative stress and diabetic cardiovascular disorders: roles of mitochondria and NADPH oxidase. *Can J Physiol Pharmacol* 2010; 88(3): 241-8.
3. Nascimento H1, Silva L, Lourenço P, Weinfurterová R, Castro E, Rego C, Ferreira H, Guerra A, Quintanilha A, Santos-Silva A, Belo L. Lipid profile in Portuguese obese children and adolescents: interaction of apolipoprotein E polymorphism with adiponectin levels. *Arch Pediatr Adolesc Med.* 2009;163(11):1030-1036.
4. Amr M. Zoair, Khaled T. Muhammad, Desoky E. Abu-Ammo, Mahmoud M. Motawea. Lipid profile and some cardiac functions in children with obesity. *Egyptian Paediatric Association Gazette* (2013) 60, 15–22.
5. Minakshi B, Chithambaram N S. Abnormalities of lipid profile in overweight and obese Indian children. *Int J Pediatr Res.*2016;3(8):584-588.doi:10.17511/ijpr.2016.8.06.
6. LaFrance S, Nolan M, Page J. et al. New hampshire childhood obesity report, foundation for healthy community; 2005.
7. Shabana T, Viswanathan V. Impact of Socio economic Status on Prevalence of Overweight and Obesity among Children and Adolescents in Urban India.*The Open Obesity Journal*,2009;1:9-14.
8. Kumar S, Mahabalaraju DK, Anuroopa MS. Prevalence of Obesity and its influencing factors among affluent school children of Davengere City.*Indian Journal of Community Medicine*, January 2007;32:15-17.
9. Freedman D, Dietz W, Srinivasan S, et al. The relation of overweight to cardiovascular risk factors among children and adolescents: the Bogalusa heart study. *Pediatrics* 1999;103:1175–82.
10. Reinhr T, Andler W, Denzer C. Cardiovascular risk factors in overweight German children and adolescents: relation to gender, age and degree of overweight. *NutrMetabCardiovasc Dis* 2005;15: 181–7.
11. Kim H, Park J, Kim H, et al. Obesity and cardiovascular risk factors in Korean children and adolescents aged 10–18 years. *Am J Epidemiol* 2006;164:787–93.
12. Chinali M, De Simone G, Roman MJ, et al. Impact of obesity on cardiac geometry and function in a population of adolescents: the strong heart study. *J Am Coll Cardiol* 2006;47: 2267–73.
13. Vitarelli A, Martino F, Capotosto L, Martino E, Colantoni C, Ashurov R, Ricci S, Conde Y, Maramao F, Vitarelli M, De Chiara S, Zanoni C. Early myocardial deformation changes in hypercholesterolemic and obese children and adolescents: a 2D and 3D speckle tracking echocardiography study. *Medicine (Baltimore)*. 2014 Sep;93(12):e71. doi: 10.1097/MD. 0000000000000071.